

## MEDICATION RESTRICTIONS PRIOR TO SURGERY OR PROCEDURES

Take no medications from the list below for one week before the procedure. These drugs can cause excessive bleeding if taken prior to surgery or biopsy. (NOTE: TYLENOL IS OK!)

Advil	Diclofenac	Measurin	Vanquish
Aggrenox	Dicumarol	Meclomen	Vitamin E
Alka Seltzer	Dipyridamole	Micranin	Voltaren
Aleve	Disalcid	Midol	Warfarin
Anacin	Doans Pills	Mobidin	Zorpin
Anaprox	Dolcin	Mobigesic	
Ansaid	Dolobid	Momentum	
Anturane	Double-A	Motrin	
APAP Fort.	Duoprin	Nabumetone	
Argesic	Duradyne	Nalfon	
Artha-G	Duragesic	Naprosyn	
Artholate	Durasal	Neocylate	
Arthropan	Dynosal	Nuprin	
Ascriptin	Ecotrin	Oxalid	
Asper Buf	Efficin	PAC	
Aspercine	Emgrin	Pabalate	
Aspergum	Empirin	Pabirin	
Aspirin	Encaprin	Panodynes	
Asproject	Equagesic	Pepto-Bismol	
Axotal	Etodolac	Percodan	
Bayer	Exedrin	Persantine	
B C Tabs	Feldene	Protension	
Buf Tabs	Fiorinal	Postel	
Buff A	Flurbiprofen	Relafen	
Buffaprin	Garlic	Rufen	
Bufferin	Gaysal-S	Sal-Favne	
Buffetts II	Gemnisyn	Salatin	
Buffinol	Ginkgo	Saletto	
Buflax	Ginseng	Sine-Off	
Butal Comp	Ibuprofen	Solocol	
Butazolidin	Indocin	Supac	
Cama	Isollyl	Synalgos	
Carpon	Kaopectate	Tandearil	
Cataflam	Ketoralac	Tenstan	
Clinoril	Lanorinal	Ticlid	
Cope	Lodine	Ticlopidine	
Coricidin	Magan	Tisma	
CP-2	Magsal	Tolectin	
Cosprin	Marnal	Toradol	
Coumadin	Major-cin	Trigesic	
Dasin	Majoral	UraceLS	

**Stop Coumadin / Warfarin five (5) days prior to the procedure. Also, check with the Anticoagulation Clinic to make sure this is OK.**