

INSTRUCTIONS FOR ALDARA CREAM APPLICATION

Kaiser Permanente, Santa Clara Medical Center
Department of Dermatology, #472
710 Lawrence Expressway
Santa Clara, CA 95051
408-851-4650

1. Aldara cream (Imiquimod) is a cream that helps your immune system to fight against several types of growths on your skin. It is used to treat warts, molluscum, sun damaged skin (actinic keratosis), and some skin cancers. Aldara causes local irritation and even ulceration when it works.
2. Aldara cream comes in little packets. Perforate the packet with a needle and squeeze just enough onto the fingertip or a Q-tip to apply thinly to the lesion and immediate surrounding areas. Only a small amount of cream is needed. You can save the rest of the medication in the packet with a ziplock bag.
3. Apply Aldara cream to the lesion at bedtime 5 times per week (M-T-W-T-F) for 6 weeks, with weekends off. The goal of the treatment is to achieve the desired local redness and mild crusting. If there is minimal redness, increase to seven times a week (M-T-W-T-F-S-S).
4. If there is too much redness or irritation, reduce accordingly. You can use as infrequently as twice a week (M-W) or three times a week (M-W-F). The goal is to obtain tolerable amount of irritation, which is the evidence that your body is trying to fight the disease.
5. If a severe local reaction occurs, such as extensive crusting or blistering, stop the Aldara cream until the reaction resolves. This usually takes 3-4 days, then restart at a reduced frequency.
6. Use the medicine for 6 to 15 weeks then stop. The duration is dependent on the frequency of the application. In general, 30 applications in total will give the best results.
7. After you stop the medication, give the skin 3 to 4 weeks to heal from the treatment. At that time, if you think the skin disease is still present, set up an appointment with me for evaluation.
8. If you have problems or questions, stop the medication and call Dermatology at 408-851-4650.